

On the $\Sigma^0 \rightarrow \Lambda^0 + \gamma$ decay in the Effective quark model with chiral $U(3) \times U(3)$ symmetry

A. N. Ivanov^{1,2,3,*†}, A. Ya. Berdnikov³, Ya. A. Berdnikov³, M. Faber¹,
V. A. Ivanova³, V. F. Kosmach³, A. V. Nikitchenko³, N. I. Troitskaya³

February 2, 2008

¹*Atominstitut der Österreichischen Universitäten, Technische Universität Wien,
Wiedner Hauptstrasse 8-10, A-1040 Wien, Österreich*

and

²*Stefan Meyer Institut für subatomare Physik, Österreichische Akademie der
Wissenschaften, Boltzmanngasse 3, A-1090, Wien, Österreich*

and

³*State Polytechnic University, Polytechnicheskaya 29,
195251, St Petersburg, Russian Federation*

Abstract

We analyse the angular distributions of the $\Sigma^0 \rightarrow \Lambda^0 \gamma$ decay rate in the laboratory and in the rest frame of the Σ^0 -hyperon in the dependence on baryon polarizations. We calculate the dynamical polarization vector of the Λ^0 -hyperon. Within the Effective quark model with chiral $U(3) \times U(3)$ symmetry (PRC **59**, 451 (1999)) we calculate the transition magnetic moment $\mu_{\Sigma^0 \Lambda^0}$. The theoretical value $\mu_{\Sigma^0 \Lambda^0} = -1.62$, measured in nuclear magnetons, agrees well with the experimental data $|\mu_{\Sigma^0 \Lambda^0}^{\text{exp}}| = (1.61 \pm 0.08)$ and the theoretical result, predicted within the naive quark model $\mu_{\Sigma^0 \Lambda^0} = (\sqrt{3}/4)(\mu_{\Sigma^-} - \mu_{\Sigma^+}) = (-1.57 \pm 0.01)$.

PACS: 13.30.Eg, 13.75.Ev, 13.88.+e, 14.20.Jn

*Corresponding author. E-mail: ivanov@kph.tuwien.ac.at, Tel.: +43-1-58801-14261, Fax: +43-1-58801-14299

†Permanent Address: State Polytechnic University, Department of Nuclear Physics, 195251 St. Petersburg, Russian Federation

1 Introduction

The Effective quark model with chiral $U(3) \times U(3)$ symmetry [1] has been successfully applied to the calculation of the non-leptonic decays of charmed Λ_c^+ -baryon [2, 3], the S-wave amplitudes of $\bar{K}N$ and K^-d scattering at threshold [5, 6] and the analysis of the polarization properties of the Λ^0 -hyperon, produced from the Quark-gluon plasma [7]. The value of the $\sigma_{\pi N}$ -term, $\sigma_{\pi N} = 60 \text{ MeV}$ calculated within the Effective quark model with chiral $U(3) \times U(3)$ symmetry [1], agrees well with the experimental value $\sigma_{\pi N}^{\text{exp}} = 61_{-4}^{+1} \text{ MeV}$, extracted from the experimental data on the energy level displacement of the ground state of pionic hydrogen [4].

In this paper we apply the Effective quark model with chiral $U(3) \times U(3)$ symmetry to the calculation of the transition magnetic moment $\mu_{\Sigma^0 \Lambda^0}$, responsible for the $\Sigma^0 \rightarrow \Lambda^0 + \gamma$ decay. A new interest to this decay is connected with the experimental search for the Kaonic Nuclear Cluster K^-pp state (the KNC $\frac{2}{K}\text{H}$) [8], predicted within the Potential model approach by Akaishi and Yamazaki [9] and the Phenomenological quantum field theory model in [10]. Experimentally the existence of the KNC $\frac{2}{K}\text{H}$ is analysed by means of the invariant-mass spectrum of the $\Lambda^0 p$ pair, induced by the decays $\frac{2}{K}\text{H} \rightarrow \Lambda^0 + p$ and $\frac{2}{K}\text{H} \rightarrow \Sigma^0 + p$. Due to the decay $\Sigma^0 \rightarrow \Lambda^0 + \gamma$ the experimental invariant-mass spectrum of the $\Lambda^0 p$ pair should contain two peaks. One of these peaks is located at the mass of the KNC $\frac{2}{K}\text{H}$. It is defined by the $\frac{2}{K}\text{H} \rightarrow \Lambda^0 + p$ decay. Another one is shifted to left from the mass of the KNC $\frac{2}{K}\text{H}$ by about 78 MeV. This shift is defined by the photon energy in the $\frac{2}{K}\text{H} \rightarrow \Sigma^0 + p \rightarrow \Lambda^0 + \gamma + p$ cascade decay.

The paper is organized as follows. In Section 2 we calculate the angular distributions of the $\Sigma^0 \rightarrow \Lambda^0 \gamma$ decay rate in the laboratory and in the rest frame of the Σ^0 -hyperon in dependence on the polarizations of the baryons. We calculate the dynamical polarization vector of the Λ^0 -hyperon. In Section 3 we calculate the transition magnetic moment $\mu_{\Sigma^0 \Lambda^0} = -1.62$, measured in nuclear magnetons. The calculation is carried out in the Effective quark model with chiral $U(3) \times U(3)$ symmetry. The theoretical result agrees well with the experimental value $|\mu_{\Sigma^0 \Lambda^0}^{\text{exp}}| = (1.61 \pm 0.08)$ [11] and the theoretical prediction within the naive quark model $\mu_{\Sigma^0 \Lambda^0} = -1.57 \pm 0.01$ [12]. In the Conclusion we discuss the obtained results.

2 Angular distribution of the $\Sigma^0 \rightarrow \Lambda^0 \gamma$ decay rate

The $\Sigma^0 \rightarrow \Lambda^0 \gamma$ decay is the $M1$ electromagnetic transition. The amplitude of this transition is defined by

$$M(\Sigma^0 \rightarrow \Lambda^0 \gamma) = \sqrt{4\pi\alpha} e_\mu^*(k, \lambda) \langle \Lambda^0(k_{\Sigma^0}, \sigma_{\Sigma^0}) | J^\mu(0) | \Sigma^0(k_{\Lambda^0}, \sigma_{\Lambda^0}) \rangle, \quad (2.1)$$

where $J^\mu(0)$ is the electromagnetic hadronic current, $e_\mu^*(k, \lambda)$ is a polarization vector of a photon and $\alpha = 1/137.036$ is the fine-structure constant. Due to the $M1$ transition the matrix element of the electromagnetic hadronic current is

$$\langle \Lambda^0(k_{\Sigma^0}, \sigma_{\Sigma^0}) | J^\mu(0) | \Sigma^0(k_{\Lambda^0}, \sigma_{\Lambda^0}) \rangle = -g_{\Sigma^0 \Lambda^0} \bar{u}_{\Lambda^0}(k_{\Lambda^0}, \sigma_{\Lambda^0}) \sigma^{\mu\nu} k_\nu u_{\Sigma^0}(k_{\Sigma^0}, \sigma_{\Sigma^0}), \quad (2.2)$$

where $\sigma^{\mu\nu} = (\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu)/2$, $g_{\Sigma^0 \Lambda^0}$ is the effective coupling constant of the transition $\Sigma^0 \rightarrow \Lambda^0$, caused by strong low-energy interactions, and $k = k_{\Sigma^0} - k_{\Lambda^0}$, and

$\bar{u}_{\Lambda^0}(k_{\Lambda^0}, \sigma_{\Lambda^0})$ and $u_{\Sigma^0}(k_{\Sigma^0}, \sigma_{\Sigma^0})$ are the bispinors of the Λ^0 and Σ^0 hyperons, normalized by $\bar{u}_Y(k_Y, \sigma_Y)u_Y(k_Y, \sigma_Y) = 2m_Y$ for $Y = \Sigma^0$ and Λ^0 .

The width of the $\Sigma^0 \rightarrow \Lambda^0 \gamma$ decay for unpolarized baryons is proportional to

$$\begin{aligned}
& \frac{1}{2} \sum_{\sigma_{\Lambda^0}=\pm 1/2} \sum_{\sigma_{\Sigma^0}=\pm 1/2} \sum_{\lambda=\pm 1} |M(\Sigma^0 \rightarrow \Lambda^0 \gamma)|^2 = \\
& = 2\pi\alpha g_{\Sigma^0 \Lambda^0}^2 \sum_{\lambda=\pm 1} e_\mu^*(k, \lambda) e_\alpha(k, \lambda) k_\nu k_\beta \text{tr}\{(m_{\Sigma^0} + \hat{k}_{\Sigma^0})\sigma^{\mu\nu}(m_{\Lambda^0} + \hat{k}_{\Lambda^0})\sigma^{\alpha\beta}\} = \\
& = 2\pi\alpha g_{\Sigma^0 \Lambda^0}^2 \sum_{\lambda=\pm 1} e_\mu^*(k, \lambda) e_\alpha(k, \lambda) k_\nu k_\beta \text{tr}\{\hat{k}_{\Sigma^0}\sigma^{\mu\nu}\hat{k}_{\Lambda^0}\sigma^{\alpha\beta}\} = 32\pi\alpha g_{\Sigma^0 \Lambda^0}^2 (k \cdot k_{\Sigma^0})(k \cdot k_{\Lambda^0}) = \\
& = 32\pi\alpha g_{\Sigma^0 \Lambda^0}^2 m_{\Sigma^0}^2 \omega_0^2,
\end{aligned} \tag{2.3}$$

where $\omega_0 = (m_{\Sigma^0}^2 - m_{\Lambda^0}^2)/2m_{\Lambda^0} = 74.5 \text{ MeV}$ [11] is the photon energy in the rest frame of the Σ^0 -hyperon.

The width of the $\Sigma^0 \rightarrow \Lambda^0 \gamma$ decay is equal to

$$\begin{aligned}
\Gamma(\Sigma^0 \rightarrow \Lambda^0 \gamma) &= \frac{\alpha}{\pi} \frac{g_{\Sigma^0 \Lambda^0}^2}{m_{\Sigma^0}} \int \delta^{(4)}(k_{\Lambda^0} + k - k_{\Sigma^0}) (k \cdot k_{\Sigma^0})^2 \frac{d^3 k_{\Lambda^0}}{E_{\Lambda^0}(\vec{k}_{\Lambda^0})} \frac{d^3 k}{\omega(\vec{k})} = \\
&= 4\alpha g_{\Sigma^0 \Lambda^0}^2 \omega_0^3.
\end{aligned} \tag{2.4}$$

Using the experimental value for the width of the $\Sigma^0 \rightarrow \Lambda^0 \gamma$ decay $\Gamma_{\text{exp}}(\Sigma^0 \rightarrow \Lambda^0 \gamma) = (8.9 \pm 0.8) \times 10^{-3} \text{ MeV}$ [11] we get $g_{\Sigma^0 \Lambda^0} = (8.6 \pm 0.4) \times 10^{-4} \text{ MeV}^{-1}$. The effective coupling constant $g_{\Sigma^0 \Lambda^0}$ can be expressed in terms of the transition magnetic moment $g_{\Sigma^0 \Lambda^0} = \mu_{\Sigma^0 \Lambda^0}/2m_N$, where $|\mu_{\Sigma^0 \Lambda^0}| = 1.61 \pm 0.08$ [11].

In the laboratory frame, where the Σ^0 -hyperon moves with a velocity $\vec{v}_{\Sigma^0} = \vec{k}_{\Sigma^0}/E_{\Sigma^0}$, the angular distribution of the $\Sigma^0 \rightarrow \Lambda^0 \gamma$ decay rate can be investigated with respect to both the relative angle $\vartheta_{\Sigma^0 \Lambda^0}$ of the $\Sigma^0 \Lambda^0$ pair and the relative angle $\vartheta_{\Sigma^0 \gamma}$ of the $\Sigma^0 \gamma$ pair.

For the unpolarized baryons the angular distribution of the Λ^0 -hyperon relative to the direction of the motion of the Σ^0 -hyperon is

$$4\pi \frac{dB(\Sigma^0 \rightarrow \Lambda^0 \gamma)}{d\Omega_{\Sigma^0 \Lambda^0}} = \frac{1}{\gamma_{\Sigma^0} \omega_0} \frac{\vec{k}_{\Lambda^0}^2 \sqrt{1 - \vec{v}_{\Sigma^0}^2}}{|\vec{k}_{\Lambda^0}| - E_{\Lambda^0} |\vec{v}_{\Sigma^0}| \cos \vartheta_{\Sigma^0 \Lambda^0}}, \tag{2.5}$$

where $\gamma_{\Sigma^0} = 1/\sqrt{1 - \vec{v}_{\Sigma^0}^2}$. The kinematical parameters of the baryons are related by $E_{\Lambda^0} - |\vec{k}_{\Lambda^0}| |\vec{v}_{\Sigma^0}| \cos \vartheta_{\Sigma^0 \Lambda^0} = (m_{\Sigma^0} - \omega_0)/\gamma_{\Sigma^0}$. This allows to obtain the energy spectrum of the Λ^0 -hyperon

$$\frac{dB(\Sigma^0 \rightarrow \Lambda^0 \gamma)}{dE_{\Lambda^0}} = \frac{1}{2\gamma_{\Sigma^0} |\vec{v}_{\Sigma^0}| \omega_0}. \tag{2.6}$$

In turn, the angular distribution of the photon relative to the direction of the motion of the Σ^0 -hyperon takes the form

$$4\pi \frac{dB(\Sigma^0 \rightarrow \Lambda^0 \gamma)}{d\Omega_{\Sigma^0 \gamma}} = \frac{1}{4\pi \gamma_{\Sigma^0}^2} \frac{1}{(1 - |\vec{v}_{\Sigma^0}| \cos \vartheta_{\Sigma^0 \gamma})^2}. \tag{2.7}$$

The photon energy in the laboratory frame is related to the photon energy in the center of mass frame as $\omega_0 = \gamma_{\Sigma^0} \omega (1 - |\vec{v}_{\Sigma^0}| \cos \vartheta_{\Sigma^0 \gamma})$. This allows to calculate the energy spectrum of the photon

$$\frac{dB(\Sigma^0 \rightarrow \Lambda^0 \gamma)}{d\omega} = \frac{1}{2\gamma_{\Sigma^0} |\vec{v}_{\Sigma^0}| \omega_0} = \frac{1}{\omega_+ - \omega_-}, \quad (2.8)$$

where $\omega_+ = \omega_0/\gamma_{\Sigma^0}(1 - |\vec{v}_{\Sigma^0}|)$ and $\omega_- = \omega_0/\gamma_{\Sigma^0}(1 + |\vec{v}_{\Sigma^0}|)$ are the maximal and minimal energy of the photon, emitted by the Σ^0 -hyperon in the laboratory frame.

In the rest frame of the Σ^0 -hyperon the angular distribution of the decay rate is equal to

$$4\pi \frac{dB(\Sigma^0 \rightarrow \Lambda^0 \gamma)}{d\Omega_{\Lambda^0}} = 1, \quad (2.9)$$

where $d\Omega_{\Lambda^0} = 2\pi \sin \vartheta_{\Lambda^0} d\vartheta_{\Lambda^0}$ and ϑ_{Λ^0} is polar angle relative to the momentum $\vec{k}_{\Lambda^0} = -\vec{k}$ of the Λ^0 -hyperon at the rest frame of the Σ^0 -hyperon and $\vartheta_\gamma = \pi - \vartheta_{\Lambda^0}$.

For the polarized hyperons, which we denote as $\vec{\Sigma}^0$ and $\vec{\Lambda}^0$, the expression (2.3) changes as follows [13] (see also [14]):

$$\begin{aligned} \sum_{\lambda=\pm 1} |M(\vec{\Sigma}^0 \rightarrow \vec{\Lambda}^0 \gamma)|^2 &= 2\pi\alpha g_{\Sigma^0 \Lambda^0}^2 \sum_{\lambda=\pm 1} e_\mu^*(k, \lambda) e_\alpha(k, \lambda) k_\nu k_\beta \\ &\times \text{tr}\{(m_{\Sigma^0} + \hat{k}_{\Sigma^0})(1 + \gamma^5 \hat{\zeta}_{\Sigma^0}) \sigma^{\mu\nu} (m_{\Lambda^0} + \hat{k}_{\Lambda^0})(1 + \gamma^5 \hat{\zeta}_{\Lambda^0}) \sigma^{\alpha\beta}\}. \end{aligned} \quad (2.10)$$

Here $\hat{\zeta}_Y = \gamma_\mu \zeta_Y^\mu$ for $Y = \Sigma^0, \Lambda^0$, where ζ_Y^μ is the 4-polarization vector of the Y -hyperon [13] (see also [14]):

$$\zeta_Y^\mu = \left(\frac{\vec{k}_Y \cdot \vec{\xi}_Y}{m_Y}, \vec{\xi}_Y + \frac{\vec{k}_Y (\vec{k}_Y \cdot \vec{\xi}_Y)}{m_Y (E_Y(\vec{k}_Y) + m_Y)} \right), \quad (2.11)$$

where $\vec{\xi}_Y$ is a 3-dimensional polarization vector of the Y -hyperon, normalized to unity $\vec{\xi}_Y^2 = 1$. The polarization vector ζ_Y^μ satisfies the constraints $\zeta_Y^2 = -1$ and $k_Y \cdot \zeta_Y = 0$.

The result of the calculation of the trace in (2.10) is

$$\begin{aligned} \sum_{\lambda=\pm 1} |M(\vec{\Sigma}^0 \rightarrow \vec{\Lambda}^0 \gamma)|^2 &= 32\pi\alpha g_{\Sigma^0 \Lambda^0}^2 \left\{ (k \cdot k_{\Sigma^0})^2 - (k \cdot k_{\Sigma^0})^2 (\zeta_{\Sigma^0} \cdot \zeta_{\Lambda^0}) \right. \\ &- m_{\Sigma^0} (m_{\Sigma^0} + m_{\Lambda^0}) (k \cdot \zeta_{\Sigma^0}) (k \cdot \zeta_{\Lambda^0}) - \frac{1}{\omega^2} (k \cdot k_{\Sigma^0})^2 (\vec{k} \times \vec{\zeta}_{\Sigma^0}) \cdot (\vec{k} \times \vec{\zeta}_{\Lambda^0}) \\ &- \frac{1}{\omega^2} (\vec{k} \times \vec{k}_{\Sigma^0}) \cdot (\vec{k} \times \vec{k}_{\Sigma^0}) (k \cdot \zeta_{\Sigma^0}) (k \cdot \zeta_{\Lambda^0}) + \frac{1}{\omega^2} (k \cdot k_{\Sigma^0}) (k \cdot \zeta_{\Lambda^0}) (\vec{k} \times \vec{k}_{\Sigma^0}) \cdot (\vec{k} \times \vec{\zeta}_{\Sigma^0}) \\ &\left. + \frac{1}{\omega^2} (k \cdot k_{\Sigma^0}) (k \cdot \zeta_{\Sigma^0}) (\vec{k} \times \vec{k}_{\Lambda^0}) \cdot (\vec{k} \times \vec{\zeta}_{\Lambda^0}) \right\}. \end{aligned} \quad (2.12)$$

We have carried out the summation over the transverse polarizations of the photon.

The angular distribution of the $\vec{\Lambda}^0$ -hyperon relative to the direction of the motion of the $\vec{\Sigma}^0$ -hyperon can be written in the following general form [13]

$$4\pi \frac{dB(\vec{\Sigma}^0 \rightarrow \vec{\Lambda}^0 \gamma)}{d\Omega_{\Sigma^0 \Lambda^0}} = 4\pi \frac{dB(\Sigma^0 \rightarrow \Lambda^0 \gamma)}{d\Omega_{\Sigma^0 \Lambda^0}} (1 + \vec{P}_{\Lambda^0} \cdot \vec{\xi}_{\Lambda^0}), \quad (2.13)$$

where $\vec{P}_{\Lambda^0}$ is a dynamical polarization vector of the Λ^0 -hyperon. Following the procedure expounded in [13] we obtain

$$\begin{aligned} \vec{P}_{\Lambda^0} = & \left\{ \frac{m_{\Sigma^0} + m_{\Lambda^0}}{m_{\Sigma^0}\omega_0^2} (k_{\Lambda^0} \cdot \zeta_{\Sigma^0}) - \frac{(\vec{k}_{\Sigma^0} \times \vec{k}_{\Lambda^0})^2}{|\vec{k}_{\Sigma^0} - \vec{k}_{\Lambda^0}|^2} \frac{k_{\Lambda^0} \cdot \zeta_{\Sigma^0}}{m_{\Sigma^0}\omega_0^2} - \frac{(\vec{k}_{\Sigma^0} \times \vec{k}_{\Lambda^0}) \cdot ((\vec{k}_{\Sigma^0} - \vec{k}_{\Lambda^0}) \times \vec{\zeta}_{\Sigma^0})}{|\vec{k}_{\Sigma^0} - \vec{k}_{\Lambda^0}|^2 m_{\Sigma^0}\omega_0} \right\} \\ & \times \left[E_{\Sigma^0} \frac{\vec{k}_{\Lambda^0}}{m_{\Lambda^0}} - \vec{k}_{\Sigma^0} - \frac{(\vec{k}_{\Sigma^0} \cdot \vec{k}_{\Lambda^0}) \vec{k}_{\Lambda^0}}{m_{\Lambda^0}(E_{\Lambda^0} + m_{\Lambda^0})} \right] - \left(\frac{\vec{k}_{\Sigma^0} \cdot \vec{\zeta}_{\Sigma^0}}{E_{\Sigma^0}} \right) \frac{\vec{k}_{\Lambda^0}}{m_{\Lambda^0}} + \frac{(\vec{k}_{\Lambda^0} \cdot \vec{\zeta}_{\Sigma^0}) \vec{k}_{\Lambda^0}}{m_{\Lambda^0}(E_{\Lambda^0} + m_{\Lambda^0})} \\ & + \frac{1}{|\vec{k}_{\Sigma^0} - \vec{k}_{\Lambda^0}|^2} \left\{ ((\vec{k}_{\Sigma^0} - \vec{k}_{\Lambda^0}) \cdot \vec{\zeta}_{\Sigma^0})(\vec{k}_{\Sigma^0} - \vec{k}_{\Lambda^0}) - \frac{(((\vec{k}_{\Sigma^0} - \vec{k}_{\Lambda^0}) \times \vec{\zeta}_{\Sigma^0}) \cdot (\vec{k}_{\Sigma^0} \times \vec{k}_{\Lambda^0})) \vec{k}_{\Lambda^0}}{m_{\Lambda^0}(E_{\Lambda^0} + m_{\Lambda^0})} \right. \\ & \left. + \frac{k_{\Lambda^0} \cdot \zeta_{\Sigma^0}}{m_{\Sigma^0}\omega_0} \left[(\vec{k}_{\Sigma^0} \times \vec{k}_{\Lambda^0}) \times (\vec{k}_{\Sigma^0} - \vec{k}_{\Lambda^0}) + \frac{(\vec{k}_{\Sigma^0} \times \vec{k}_{\Lambda^0})^2 \vec{k}_{\Lambda^0}}{m_{\Lambda^0}(E_{\Lambda^0} + m_{\Lambda^0})} \right] \right\}. \end{aligned} \quad (2.14)$$

In the rest frame of the $\vec{\Sigma}^0$ -hyperon the dynamical polarization vector of the $\vec{\Lambda}^0$ -hyperon takes the form

$$\vec{P}_{\Lambda^0} = -\frac{m_{\Sigma^0}}{m_{\Lambda^0}} (\vec{n}_{\Lambda^0} \cdot \vec{\xi}_{\Sigma^0}) \vec{n}_{\Lambda^0}, \quad (2.15)$$

where $\vec{n}_{\Lambda^0} = \vec{k}_{\Lambda^0}/\omega_0$. Since in the rest frame of the $\vec{\Sigma}^0$ -hyperon $\vec{k}_{\Lambda^0} = -\vec{k}$, $\vec{n}_{\Lambda^0}$ is a unit vector $\vec{n}_{\Lambda^0}^2 = 1$.

The dynamical polarization of the $\vec{\Lambda}^0$ -hyperon, averaged over all directions of the vector $\vec{n}_{\Lambda^0}$, amounts to

$$\langle \vec{P}_{\Lambda^0} \rangle = -\frac{1}{3} \frac{m_{\Sigma^0}}{m_{\Lambda^0}} \vec{\xi}_{\Sigma^0}. \quad (2.16)$$

Our results (2.15) and (2.16) agree well with those obtained in [15]–[17].

3 Transition magnetic moment $\mu_{\Sigma^0\Lambda^0}$

The transition magnetic moment $\mu_{\Sigma^0\Lambda^0}$ responsible for the $\Sigma^0 \rightarrow \Lambda^0 + \gamma$ decay is related to the effective coupling constant $g_{\Sigma^0\Lambda^0}$ by $\mu_{\Sigma^0\Lambda^0} = 2m_N g_{\Sigma^0\Lambda^0}$. The calculation of the effective coupling constant $g_{\Sigma^0\Lambda^0}$ we carry out in the Effective quark model with chiral $U(3) \times U(3)$ symmetry [1]–[7]. For this aim we calculate the matrix element of the electromagnetic hadronic current $\langle \Lambda^0(k_{\Sigma^0}, \sigma_{\Sigma^0}) | J^\mu(0) | \Sigma^0(k_{\Lambda^0}, \sigma_{\Lambda^0}) \rangle$.

After the application the reduction technique and the equations of motion [1, 2] the matrix element $\langle \Lambda^0(k_{\Sigma^0}, \sigma_{\Sigma^0}) | J^\mu(0) | \Sigma^0(k_{\Lambda^0}, \sigma_{\Lambda^0}) \rangle$ takes the form

$$\begin{aligned} \langle \Lambda^0(k_{\Sigma^0}, \sigma_{\Sigma^0}) | J^\mu(0) | \Sigma^0(k_{\Lambda^0}, \sigma_{\Lambda^0}) \rangle = & -\frac{1}{2} g_B^2 \int d^4x d^4y e^{ik_{\Lambda^0} \cdot x - ik_{\Sigma^0} \cdot y} \\ & \times \bar{u}_{\Lambda^0}(k_{\Lambda^0}, \sigma_{\Lambda^0}) \langle 0 | T(\eta_{\Lambda^0}(x) J_\mu(0) \bar{\eta}_{\Sigma^0}(y)) | 0 \rangle_c u_{\Sigma^0}(k_{\Sigma^0}, \sigma_{\Sigma^0}), \end{aligned} \quad (3.1)$$

where T is a time-ordering operator, the index c in $\langle 0 | \dots | 0 \rangle_c$ means the calculation of the *connected* part of the vacuum expectation value, $J_\mu(0)$ is the electromagnetic quark current

$$J_\mu(0) = \frac{2}{3} \bar{u}_\ell(0) \gamma_\mu u_\ell(0) - \frac{1}{3} \bar{d}_\ell(0) \gamma_\mu d_\ell(0) - \frac{1}{3} \bar{s}_\ell(0) \gamma_\mu s_\ell(0), \quad (3.2)$$




Figure 1: Feynman diagrams of the $\Sigma^0 \rightarrow \Lambda^0$ transition in the Effective quark model with chiral $U(3) \times U(3)$ symmetry.

where $u_\ell(0)$, $d_\ell(0)$ and $s_\ell(0)$ are current quark fields. The summation over the “colour” index $\ell = 1, 2, 3$ is assumed. Then, $\eta_{\Lambda^0}(x)$ and $\bar{\eta}_{\Sigma^0}(y)$ are the three-quark densities [1, 2, 18]

$$\begin{aligned}\eta_{\Lambda^0}(x) &= -\sqrt{\frac{2}{3}} \varepsilon^{ijk} \{ [\bar{u}_i^c(x) \gamma_\mu s_j(x)] \gamma^\mu \gamma^5 d_k(x) - [\bar{d}_i^c(x) \gamma_\mu s_j(x)] \gamma^\mu \gamma^5 u_k(x) \}, \\ \eta_{\Sigma^0}(x) &= -\sqrt{\frac{1}{2}} \varepsilon^{ijk} \{ [\bar{u}_i^c(x) \gamma_\mu d_j(x)] \gamma^\mu \gamma^5 s_k(x) + [\bar{d}_i^c(x) \gamma_\mu u_j(x)] \gamma^\mu \gamma^5 s_k(x) \} \quad (3.3)\end{aligned}$$

coupled to the Λ^0 and Σ^0 hyperons [1, 2]

$$\mathcal{L}_{\text{eff}}(x) = \frac{g_B}{\sqrt{2}} \bar{\Lambda}^0(x) \eta_{\Lambda^0}(x) + \frac{g_B}{\sqrt{2}} \bar{\Sigma}^0(x) \eta_{\Sigma^0}(x) + \text{h.c.} \quad (3.4)$$

The three-quark density $\bar{\eta}_{\Sigma^0}(x)$ is equal to

$$\bar{\eta}_{\Sigma^0}(x) = +\sqrt{\frac{1}{2}} \varepsilon^{ijk} \{ \bar{s}_i(x) \gamma^\mu \gamma^5 [\bar{d}_j(x) \gamma_\mu u_k^c(x)] + \bar{s}_i(x) \gamma^\mu \gamma^5 [\bar{u}_j(x) \gamma_\mu d_k^c(x)] \}. \quad (3.5)$$

The coupling constant g_B is calculated in [1, 19]: $g_B = 1.37 \times 10^{-4} \text{ MeV}^{-2}$.

The matrix element (3.1) is defined by Feynman diagrams in Fig.1. In the coordinate representation the analytical expression of the matrix element (3.1) takes the form

$$\begin{aligned}\langle \Lambda^0(k_{\Sigma^0}, \sigma_{\Sigma^0}) | J^\mu(0) | \Sigma^0(k_{\Lambda^0}, \sigma_{\Lambda^0}) \rangle &= -\frac{\sqrt{3}}{2} g_B^2 \int d^4x d^4y e^{ik_{\Lambda^0} \cdot x - ik_{\Sigma^0} \cdot y} \\ &\times \bar{u}_{\Lambda^0}(k_{\Lambda^0}, \sigma_{\Lambda^0}) \gamma^\alpha \gamma^5 S_F^{(d)}(x-y) \gamma_\beta S_F^{(u)}(y) \gamma_\mu S_F^{(u)}(-x) \gamma_\alpha S_F^{(s)}(x-y) \gamma^\beta \gamma^5 u_{\Sigma^0}(k_{\Sigma^0}, \sigma_{\Sigma^0}) \\ &- \frac{\sqrt{3}}{2} g_B^2 \int d^4x d^4y e^{ik_{\Lambda^0} \cdot x - ik_{\Sigma^0} \cdot y} \text{tr} \{ \gamma_\alpha S_F^{(s)}(x-y) \gamma_\beta S_F^{(d)}(y-x) \}\end{aligned}$$

$$\begin{aligned}
& \times \bar{u}_{\Lambda^0}(k_{\Lambda^0}, \sigma_{\Lambda^0}) \gamma^\alpha \gamma^5 S_F^{(u)}(x) \gamma_\mu S_F^{(u)}(-y) \gamma^\beta \gamma^5 u_{\Sigma^0}(k_{\Sigma^0}, \sigma_{\Sigma^0}) \\
& - \frac{\sqrt{3}}{2} g_B^2 \int d^4x d^4y e^{ik_{\Lambda^0} \cdot x - ik_{\Sigma^0} \cdot y} \\
& \times \bar{u}_{\Lambda^0}(k_{\Lambda^0}, \sigma_{\Lambda^0}) \gamma^\alpha \gamma^5 S_F^{(u)}(x-y) \gamma_\beta S_F^{(d)}(y) \gamma_\mu S_F^{(d)}(-x) \gamma_\alpha S_F^{(s)}(x-y) \gamma^\beta \gamma^5 u_{\Sigma^0}(k_{\Sigma^0}, \sigma_{\Sigma^0}) \\
& - \frac{\sqrt{3}}{2} g_B^2 \int d^4x d^4y e^{ik_{\Lambda^0} \cdot x - ik_{\Sigma^0} \cdot y} \text{tr}\{\gamma_\alpha S_F^{(s)}(x-y) \gamma_\beta S_F^{(u)}(y-x)\} \\
& \times \bar{u}_{\Lambda^0}(k_{\Lambda^0}, \sigma_{\Lambda^0}) \gamma^\alpha \gamma^5 S_F^{(d)}(x) \gamma_\mu S_F^{(d)}(-y) \gamma^\beta \gamma^5 u_{\Sigma^0}(k_{\Sigma^0}, \sigma_{\Sigma^0}).
\end{aligned} \tag{3.6}$$

We have taken into account that only the isovector part $(\bar{u}_\ell(0)\gamma_\mu u_\ell(0) - \bar{d}_\ell(0)\gamma_\mu d_\ell(0))/2$ of the electromagnetic current gives the contribution to the transition $\Sigma^0 \rightarrow \Lambda^0$.

In the momentum representation Eq.(3.6) reduces to the form

$$\begin{aligned}
\langle \Lambda^0(k_{\Sigma^0}, \sigma_{\Sigma^0}) | J^\mu(0) | \Sigma^0(k_{\Lambda^0}, \sigma_{\Lambda^0}) \rangle &= \frac{\sqrt{3}}{2} g_B^2 \bar{u}_{\Lambda^0}(k_{\Lambda^0}, \sigma_{\Lambda^0}) \int \frac{d^4k_1}{(2\pi)^4 i} \frac{d^4k_2}{(2\pi)^4 i} \\
& \times \left[\gamma^\alpha \gamma^5 \frac{1}{m_d - \hat{k}_2} \gamma_\beta \frac{1}{m_u - \hat{k}_1} \gamma_\mu \frac{1}{m_u - \hat{k}_{\Sigma^0} + \hat{k}_{\Lambda^0} - \hat{k}_1} \gamma_\alpha \frac{1}{m_s - \hat{k}_{\Sigma^0} - \hat{k}_1 + \hat{k}_2} \gamma^\beta \gamma^5 \right. \\
& + \text{tr} \left\{ \gamma_\beta \frac{1}{m_d - \hat{k}_2} \gamma_\alpha \frac{1}{m_s - \hat{k}_2 - \hat{k}_{\Lambda^0} + \hat{k}_1} \right\} \gamma^\alpha \gamma^5 \frac{1}{m_u - \hat{k}_1} \gamma_\mu \frac{1}{m_u - \hat{k}_{\Sigma^0} + \hat{k}_{\Lambda^0} - \hat{k}_1} \gamma^\beta \gamma^5 \\
& + \gamma^\alpha \gamma^5 \frac{1}{m_u - \hat{k}_2} \gamma_\beta \frac{1}{m_d - \hat{k}_1} \gamma_\mu \frac{1}{m_d - \hat{k}_{\Sigma^0} + \hat{k}_{\Lambda^0} - \hat{k}_1} \gamma_\alpha \frac{1}{m_s - \hat{k}_{\Sigma^0} - \hat{k}_1 + \hat{k}_2} \gamma^\beta \gamma^5 \\
& \left. + \text{tr} \left\{ \gamma_\beta \frac{1}{m_u - \hat{k}_2} \gamma_\alpha \frac{1}{m_s - \hat{k}_2 - \hat{k}_{\Lambda^0} + \hat{k}_1} \right\} \gamma^\alpha \gamma^5 \frac{1}{m_d - \hat{k}_1} \gamma_\mu \frac{1}{m_d - \hat{k}_{\Sigma^0} + \hat{k}_{\Lambda^0} - \hat{k}_1} \gamma^\beta \gamma^5 \right] \\
& \times u_{\Sigma^0}(k_{\Sigma^0}, \sigma_{\Sigma^0}).
\end{aligned} \tag{3.7}$$

Applying the heavy-baryon technique, accepted for the analysis of baryon exchanges within Chiral Perturbation Theory (ChPT) [20] (see also [21, 22] and [23]), to the calculation of the momentum integrals [1] we get (see also [6]):

$$g_{\Sigma^0 \Lambda^0} = \frac{g_B^2}{\sqrt{3}} \frac{\langle \bar{q}q \rangle}{m_N} \frac{m}{16\pi^2} \ln \left(1 + \frac{\Lambda_\chi^2}{m^2} \right) = -8.62 \times 10^{-4} \text{ MeV}^{-4}, \tag{3.8}$$

where $\langle \bar{q}q \rangle$ is the quark condensate [1, 24]:

$$\langle \bar{q}q \rangle = -\frac{3m}{4\pi^2} \left[\Lambda_\chi^2 - m^2 \ln \left(1 + \frac{\Lambda_\chi^2}{m^2} \right) \right] = (-253 \text{ MeV})^3, \tag{3.9}$$

calculated at the normalization scale $\mu = \Lambda_\chi$, where $\Lambda_\chi = 940 \text{ MeV}$ is the scale of the spontaneous breaking of chiral symmetry [1], and for $m_u = m_d = m = 330 \text{ MeV}$, the constituent quark mass defined in the chiral limit [1, 24]. For the calculation of the integrals we have neglected the mass differences between Σ^0 , Λ^0 and N and have set $m_{\Sigma^0} = m_{\Lambda^0} = m_N$. Hence, the accuracy of the theoretical value (3.8) is not better than 20 %.

For the effective coupling constant (3.8) the transition magnetic moment $\mu_{\Sigma^0 \Lambda^0}$ is equal to

$$\mu_{\Sigma^0 \Lambda^0} = 2m_N g_{\Sigma^0 \Lambda^0} = \frac{g_B^2}{\sqrt{3}} \langle \bar{q}q \rangle \frac{m}{8\pi^2} \ln \left(1 + \frac{\Lambda_\chi^2}{m^2} \right) = -1.62, \tag{3.10}$$

which agrees well with both the experimental data $|\mu_{\Sigma^0\Lambda^0}^{\text{exp}}| = (1.61 \pm 0.08)$ and the theoretical value $\mu_{\Sigma^0\Lambda^0} = (\sqrt{3}/4)(\mu_{\Sigma^-} - \mu_{\Sigma^+}) = (-1.57 \pm 0.01)$, obtained within the naive quark model in terms of the magnetic moments of the charged Σ -hyperons $\mu_{\Sigma^-} = (-1.160 \pm 0.025)$ and $\mu_{\Sigma^+} = (2.458 \pm 0.010)$ [11]. In the naive quark model the magnetic moments μ_{Σ^-} and μ_{Σ^+} are defined in terms of the magnetic moments of constituent quarks $\mu_{\Sigma^-} = (4\mu_d - \mu_s)/3$ and $\mu_{\Sigma^+} = (4\mu_u - \mu_s)/3$. For the transition magnetic moment $\mu_{\Sigma^0\Lambda^0}$ this gives $\mu_{\Sigma^0\Lambda^0} = (\mu_d - \mu_u)/\sqrt{3}$ [12].

As a result of the calculation of the momentum integrals in Eq.(3.7) there appears the term proportional to $\bar{u}_{\Lambda^0}\gamma^\mu u_{\Sigma^0}$. Since in the heavy-baryon limit the spatial part of this current $\bar{u}_{\Lambda^0}\vec{\gamma}u_{\Sigma^0}$ is of order of $O(1/m_N^2)$, in the heavy-baryon limit it is equivalent to zero relative to the contribution proportional to $\bar{u}_{\Lambda^0}\sigma^{\mu\nu}k_\nu u_{\Sigma^0}$. In turn, the time-component $\bar{u}_{\Lambda^0}\gamma^0 u_{\Sigma^0}$ vanishes due to the orthogonality of the spinorial functions $\varphi_{\Lambda^0}^\dagger\varphi_{\Sigma^0} = 0$.

4 Conclusion

We have analysed the angular distribution of the Λ^0 -hyperons, produced in the $\Sigma^0 \rightarrow \Lambda^0\gamma$ decay, both in the laboratory and in the rest frame of the Σ^0 -hyperon in dependence on the polarizations of baryons. We have calculated the dynamical polarization vector of the Λ^0 -hyperon. The obtained results can be used for the analysis of the polarization properties of the Σ^0 -hyperons, produced in high-energy nucleus-nucleus collisions and in the ${}^2_K\text{H} \rightarrow \Sigma^0 p$ decays of the Kaonic Nuclear Cluster ${}^2_K\text{H}$ (the deeply bound K^-pp state). A measurement of the polarization of the Σ^0 -hyperon can be carried out through the measurement of the polarization of the Λ^0 -hyperon.

The partial width of the $\Sigma^0 \rightarrow \Lambda^0\gamma$ decay is defined by the transition magnetic moment $\mu_{\Sigma^0\Lambda^0}$. The calculation of $\mu_{\Sigma^0\Lambda^0}$ we have carried out within the Effective quark model with chiral $U(3) \times U(3)$ symmetry. The theoretical value $\mu_{\Sigma^0\Lambda^0} = -1.62$ agrees well with both the experimental one $|\mu_{\Sigma^0\Lambda^0}^{\text{exp}}| = (1.61 \pm 0.08)$ and the predicted within the naive quark model $\mu_{\Sigma^0\Lambda^0} = (\sqrt{3}/4)(\mu_{\Sigma^-} - \mu_{\Sigma^+}) = (-1.57 \pm 0.01)$.

In our approach the transition magnetic moment $\mu_{\Sigma^0\Lambda^0}$ does not depend on the mass of constituent s -quark. This agrees well with the naive quark model, where $\mu_{\Sigma^0\Lambda^0}$ is defined by the magnetic moments of the constituent u - and d -quarks only: $\mu_{\Sigma^0\Lambda^0} = (\sqrt{3}/4)(\mu_{\Sigma^-} - \mu_{\Sigma^+}) = (\mu_d - \mu_u)/\sqrt{3}$ [12].

References

- [1] A. N. Ivanov, M. Nagy, and N. I. Troitskaya, Phys. Rev. **C59**, 451 (1999).
- [2] Ya. A. Berdnikov, A. N. Ivanov, V. F. Kosmach, and N.I. Troitskaya, Phys. Rev. C **60**, 015201 (1999).
- [3] A. Ya. Berdnikov, Ya. A. Berdnikov, A. N. Ivanov, V. F. Kosmach, M. D. Scadron, and N. I. Troitskaya, Phys. Rev. D **64**, 014027 (2001).
- [4] H.-Ch. Schröder *et al.*, Eur. Phys. J. C **21**, 473 (2001).

- [5] A. N. Ivanov, M. Cargnelli, M. Faber, J. Marton, N. I. Troitskaya, and J. Zmeskal, Eur. Phys. J. A **21**, 11 (2004); nucl-th/0310081.
- [6] A. N. Ivanov, M. Cargnelli, M. Faber, H. Fuhrmann, V. A. Ivanova, J. Marton, N. I. Troitskaya, and J. Zmeskal, Eur. Phys. J. A **23**, 79 (2005); nucl-th/0406053.
- [7] A.Ya. Berdnikov, Ya. A. Berdnikov, A. N. Ivanov, V. A. Ivanova, V. F. Kosmach, V. M. Samsonov, N. I. Troitskaya, and V. Thatar Vento, Acta Phys. Hung. A **22**, 139 (2005), nucl-th/0312045. A. Andronic *et al.*,
- [8] it Measurements of Strange Baryons and Kaonic Nuclear Clusters with FOPI; T. Yamazaki, K. Suzuki, L. Fabietti, P. Kienle, and N. Hermann, *Search for K Nuclear Clusters in Nuclear Collisions with FOPI*, (unpublished) 2005; M. Agnello *et al.* (the FINUDA Collaboration), Phys. Rev. Lett. **94**, 212303 (2005).
- [9] Y. Akaishi and T. Yamazaki, Phys. Rev. C **65**, 044005 (2002); T. Yamazaki and Y. Akaishi, Phys. Lett. B **535**, 70 (2002).
- [10] A. N. Ivanov, P. Kienle, J. Marton, and E. Widmann, *Phenomenological model of the Kaonic Nuclear Cluster K^-pp in the ground state*, nucl-th/0512037.
- [11] S. Eidelman *et al.* (The Particle Data Group), Phys. Lett. B **595**, 1 (2004).
- [12] (see [11] p.922)
- [13] L. L. Landau und E. M. Lifschitz, in *LEHRBUCH DER THEORETISCHEN PHYSIK*, Band IVa, *RELATIVISTISCHE QUANTENTHEORIE* von W. B. Berestet-zki, E. M. Lifschitz, und L. P. Pitajewski, Akademie-Verlag, Berlin, 1971, ss. 111–116, 270–274.
- [14] A. Ya. Berdnikov, Ya. A. Berdnikov, A. N. Ivanov, V. A. Ivanova, V. F. Kosmach, and N. I. Troitskaya, Eur. Phys. J. A **12**, 341 (2001).
- [15] R. Gatto, Phys. Rev. **109**, 610 (1958).
- [16] G. Feldman and T. Fulton, Nucl. Phys. **8**, 106 (1958).
- [17] N. Byers and H. Burkhardt, Phys. Rev. **121**, 281 (1961).
- [18] Sni-lin Zhu, W-Y. P. Hwang, and Ze-sen Yang, Phys. Rev. D **57**, 1527 (1998) and references therein.
- [19] The coupling constant g_B is defined by [1]:

$$g_{\pi NN} = g_B^2 \frac{2}{3} \frac{m}{F_\pi} \frac{\langle \bar{q}q \rangle^2}{m_N^2},$$

where $g_{\pi NN} = 13.21$ is the πNN coupling constant, $F_\pi = 92.4 \text{ MeV}$ is the PCAC constant of π -meson, $\langle \bar{q}q \rangle = (-253)^3 \text{ MeV}^3$ is the quark condensate (3.9), and $m = 330 \text{ MeV}$ is the constituent quark mass, calculated in the chiral limit. This results in $g_B = 1.37 \times 10^{-4} \text{ MeV}^{-2}$.

- [20] J. Gasser, Nucl. Phys. Proc. Suppl. **86**, 257 (2000) and references therein. H. Leutwyler, PiN Newslett. **15**, 1 (1999); B. Borasoy and B. Holstein, Eur. Phys. J. C **6**, 85 (1999); Phys. Rev. D **59**, 094025 (1999); Phys. Rev. D **60**, 054021 (1999); Ulf-G. Meißner, PiN Newslett. **13**, 7 (1997); H. Leutwyler, Ann. of Phys. **235**, 165 (1994); G. Ecker, Prog. Part. Nucl. Phys. **36**, 71 (1996); Prog. Part. Nucl. Phys. **35**, 1 (1995); Nucl. Phys. Proc. Suppl. **16**, 581 (1990); E. Jenkins and A. V. Manohar, Phys. Lett. B **225**, 558 (1991); J. Gasser, Nucl. Phys. B **279**, 65 (1987); J. Gasser and H. Leutwyler, Nucl. Phys. B **250**, 465 (1985); Ann. of Phys. **158**, 142 (1984); Phys. Lett. B **125**, 321 (1983).
- [21] E. Witten, Nucl. Phys. B **160**, 57 (1979), In the multicolour QCD with gauge $SU(N_C)$ symmetry, where N_C is the number of quark degrees of freedom, baryon masses are proportional to N_C at $N_C \rightarrow \infty$. Hence, a heavy-baryon limit can be identified with a leading order in the large N_C expansion (see also R. Flores-Mendieta, *Topics on heavy-baryon ChPT in the large N_C limit*, hep-ph/0207138 and references therein).
- [22] A. N. Ivanov, N. I. Troitskaya, H. Oberhummer, and M. Faber, Eur. Phys. J. A **7**, 519 (2000), Eur. Phys. J. A **8**, 129 (2000), Eur. Phys. J. A **8**, 223 (2000); A. N. Ivanov, V. A. Ivanova, H. Oberhummer, N. I. Troitskaya, and M. Faber, Eur. Phys. J. A **12**, 87 (2001); A. N. Ivanov, M. Cargnelli, M. Faber, H. Fuhrmann, V. A. Ivanova, J. Marton, N. I. Troitskaya, and J. Zmeskal, Eur. Phys. J. A **23**, 1 (2005); nucl-th/0406081.
- [23] T. E. O. Ericson and A. N. Ivanov, *Isospin breaking in threshold πN scattering by radiative processes*, hep-ph/0503277.
- [24] A. N. Ivanov, M. Nagy and N. I. Troitskaya, Intern. J. Mod. Phys. A **7**, 7305 (1992); A. N. Ivanov, Intern. J. Mod. Phys. A **8**, 853 (1993); A. N. Ivanov, N. I. Troitskaya and M. Nagy, Intern. J. Mod. Phys. A **8**, 2027, 3425 (1993).